



LIQUID NITROGEN TREATMENT & AFTERCARE

Liquid nitrogen is often used to treat skin lesions such as warts, actinic keratoses, and seborrheic keratoses. Freezing damages the treated lesion and helps prevent the continuous growth of that lesion. Some lesions may require multiple treatments.

What to expect

- Liquid Nitrogen is applied to the top of skin lesion and can extend to the surrounding tissue. Freezing can cause stinging and burning pain.
- Within minutes the treated lesion may become red and begin to swell.
- A blister may form, often there is a small amount of bleeding into the blister, and it might turn dark purple or black.
- The blister will flatten and slough off in 2-4 weeks. Healing times will vary by location.

How to care for the treated area

- As the blister may be tender for the first few days, you may bandage the area. However, most often it does not require a bandage.
- Do not submerge treated areas (swimming, washing dishes, hot tubs) during the first 24hours.
- Gently clean the area daily, with mild soap. Ok to apply Vaseline to area.
- If the area is excessively uncomfortable, you may take over-the-counter pain relievers such as Tylenol or Ibuprofen.
- **Follow up-** If the lesion is still present after 30 days, please schedule an appointment. Again, some lesions will require multiple treatments.

SILVERDALE

3505 NW Anderson Hill Rd
Suite 201 · Silverdale, WA 98383

360-698-6859

GIG HARBOR

4700 Point Fosdick Dr
Suite 219 · Gig Harbor, WA 98335

253-851-7733

TACOMA

4707 South 19th Street
Suite 130 · Tacoma, WA 98405

253-752-2273